

# EXPERIMENTAL DETERMINATION OF NMR TENSORS UTILIZING LCNMR SPECTROSCOPY AND COMPARISON OF RESULTS WITH THE COMPUTED ONES

Jukka Jokisaari

NMR Research Group, Department of Physical Sciences, P.O.Box 3000, FIN-90014 University of Oulu, Finland, Email:Jukka.Jokisaari@oulu.fi

NMR spectroscopy of molecules dissolved in liquid crystals (LCNMR) is a powerful experimental means to derive elements of the nuclear shielding, indirect spin-spin coupling and quadrupole coupling tensors. This is due to the fact that the anisotropic interactions between liquid crystal and solute molecules lead to partial orientation of solute molecules which in turn causes anisotropic contributions to parameters resolvable from experimental NMR spectra. In isotropic environments, only averages of the respective tensors may be determined. In order to get accurate tensorial properties it is necessary, in most cases, to derive the molecular structure and orientational order parameters so completely as possible through the direct dipole-dipole couplings. This in turn necessitates the correction of the experimental dipolar couplings for vibrational and reorientational motions.

A general introduction to LCNMR will be given. Experimental results will be compared with the computed ones for some selected cases from the past (1-3) and the present (4-7).

## References

1. J. Jokisaari, K. Räisänen, L. Lajunen, A. Passoja and P. Pyykkö, Proton, carbon and cadmium NMR measurements and relativistic calculation of the cadmium-carbon coupling tensor in dimethyl cadmium, *J. Magn. Reson.* **31**, 121 (1978).
2. J. Jokisaari, K. Räisänen, J. Kuonanoja, P. Pyykkö and L. Lajunen, The  $r_{\alpha}$ -structure and anisotropy of the Hg-C indirect coupling constant of merthylmercury nitrate dissolved in nematic and lyotropic liquid crystals, *Molec. Phys.* **39**, 715 (1980).
3. P. Diehl, J. Jokisaari, J. Amrein, T. Väänänen and P. Pyykkö, Determination of the  $^{13}\text{C}$ - $^{13}\text{C}$  and  $^{13}\text{C}$ - $^{15}\text{N}$  coupling constant anisotropies of acetonitrile partially oriented by nematic liquid crystals, *J. Magn. Reson.* **48**, 495 (1982).
4. J. Kaski, J. Vaara and J. Jokisaari,  $^{13}\text{C}$ - $^{13}\text{C}$  spin-spin coupling tensors in benzene as determined experimentally by liquid crystal NMR and theoretically by *ab initio* calculations, *J. Am. Chem. Soc.*, **118**, 8879-8886 (1996).
5. J. Kaski, P. Lantto, J. Vaara and J. Jokisaari, "Experimental and theoretical *ab initio* study of the  $^{13}\text{C}$ - $^{13}\text{C}$  spin-spin coupling and  $^1\text{H}$  and  $^{13}\text{C}$  shielding tensors in ethane, ethene and ethyne", *J. Am. Chem. Soc.* **120**, 3993-4005 (1998).
6. J. Vaara, J. Kaski and J. Jokisaari, Indirect fluorine coupling anisotropies in p-difluorobenzene: Implications to orientation and structure determination of fluorinated liquid crystals, *J. Phys. Chem A*, **103**, 5675-5684 (1999).
7. P. Lantto, J. Kaski, J. Vaara and J. Jokisaari, Spin-spin coupling tensors in fluoromethanes, *Chemistry-A European Journal*, **6**, 1395-1406 (2000).